

Amino Acid Balancing in the Context of MP and RUP Requirements

Charles G. Schwab, Ryan S. Ordway, and Nancy L. Whitehouse

Department of Animal and Nutritional Sciences

University of New Hampshire

Durham, NH 03824

Introduction

The protein requirements of lactating dairy cows continue to be refined. In NRC (1971, 1978), dietary requirements were expressed as CP and metabolic requirements as digestible protein. In NRC (1989), dietary requirements were expressed as CP or degraded intake CP (DIP) and undegraded intake CP (UIP), and metabolic requirements as absorbed protein. Mean values of ruminal degradability for common feeds, derived from in vivo and in situ studies using sheep and cattle, were reported. A fixed intestinal digestibility of 80% for RUP and microbial true protein was used for predicting passage of absorbed protein. In NRC (2001), dietary requirements are expressed as rumen-degradable CP (RDP) and rumen-undegradable CP (RUP) and metabolic requirements are expressed as metabolizable protein (MP). In contrast to NRC (1989): 1) microbial CP flows are predicted from intake of total digestible OM intake instead of NE intake, 2) a mechanistic system is used for predicting the RDP and RUP content of feeds that recognizes that the proportional content of these two fractions is not constant and is affected by DM intake and diet composition, 3) variable estimates of digestibility are assigned to the RUP fraction of each feed, and 4) flows of digestible essential amino acids (EAA) and their content in MP are predicted. Amino acid requirements were not established, but dose-response curves that relate measured milk protein content and yield responses to changes of predicted percentages of lysine (Lys) and methionine (Met) in MP are provided.

The purpose of this paper is to emphasize the importance of providing the lactating cow with the correct balance of AA in MP, to explain why some AA are more limiting than others, to review how one balances for Lys and Met in the context of balancing for MP and RUP and some production responses that have occurred when the practice is implemented, and to share some preliminary data aimed at extending the application of NRC (2001) to predict lactation responses from changes in supply of MP-Lys and MP-Met.

Why AA Balancing is Important

Absorbed AA, not protein per se, are the required nutrients. In ruminants, absorbed AA are provided by ruminally-synthesized bacteria, RUP, and endogenous protein. Amino acids are the building blocks for tissue and milk proteins. They are joined together in proteins according to a predetermined genetic code. Therefore, the AA composition of a protein is the same every time

it is synthesized. Hence, the AA composition of synthesized proteins is not affected by the amount or profile of absorbed AA.

While the profile of absorbed AA does not affect the AA composition of synthesized proteins, the profile of absorbed EAA does affect the amount of protein that can be synthesized. This is widely documented in swine (NRC, 1998) and poultry (NRC, 1994). When EAA are absorbed in the profile as required by the animal, their efficiency of use for protein synthesis is maximized and the requirement for total absorbed AA is reduced to a minimum. As a result, catabolism of “left-over” AA is minimal and urinary N excretion is lessened. In contrast, efficiency of use of AA for protein synthesis is less than maximal when the profile of absorbed EAA is less than ideal. In this case, it's the supply of the first-limiting EAA that determines the extent of protein synthesis, not MP (total AA) supply.

Limiting Amino Acids

Limiting AA are those that are in shortest supply relative to requirements. Methionine (Met), lysine (Lys), and histidine (His) have been identified most often as the most limiting AA for lactating dairy cows.

As reviewed in NRC (2001), Met is typically first limiting when most of the RUP is provided by soybean protein, animal-derived proteins, or a combination of the two. This is because of their low concentrations of Met as compared to milk and bacterial protein. As shown in Table 1, Met in milk and bacteria are 2.6-2.7% of CP; in soybean, blood, feather, and meat meals, Met is only 0.8-1.4% of CP. In contrast, Lys is first-limiting when corn and feeds of corn origin provide most or all of RUP (NRC, 2001). Again, these findings are not surprising. Lysine in milk and bacteria are 7.6 and 7.9% of CP, respectively, whereas in corn silage, corn, corn distillers, and corn gluten meal, Lys is only 1.7-2.8% of CP.

As might be expected from the data presented in Table 1, Met and Lys have been identified as co-limiting AA for milk protein production when cows were fed corn silage-based diets containing complementary feed proteins (NRC, 2001). This is because only a small number of feeds have concentrations of either Lys or Met in CP that is as high as the concentrations observed in milk and bacterial protein. Thus, while several feeds are complementary to each other [e.g., corn (low Lys, high Met) and soybean meal (high Lys, low Met)], their combined use cannot be expected to eliminate Met and Lys as the most limiting AA. This is especially when corn-based diets are fed.

More recently, His has been shown to be more limiting than Lys or Met when cows are fed grass silage based diets (Kim et al., 1999, 2000, 2001a, 2001b; Huhtanen et al., 2002; Korhonen et al., 2000; Vanhatalo et al., 1999). In all cases, the diets were devoid of corn and contained barley and oats as the supplemental

energy feeds. The diets were fed with or without feather meal as the sole source of supplemental RUP. While these diets are not common in the United States, these findings make an important contribution to our understanding of AA requirements and highlight the impact that diet composition has on the sequence of AA limitation. Why did the cows respond to abomasal and intravenous infusions of His?

One cannot be certain as to why His was more limiting than Lys and Met, but the data presented in Table 1 gives at least two clues. First, His may be more limiting in ruminally synthesized bacteria than either Lys or Met for milk protein synthesis. As noted in Table 1, His is 2.0% of CP in rumen bacteria and His is 2.7% of CP in milk. In contrast, concentrations of Lys and Met are both very similar in rumen bacteria and milk (7.9 and 7.6%, and 2.6 and 2.7%, respectively). This is mentioned because in these experiments, it is expected that bacterial protein constituted a larger percentage of total MP than in cows fed corn-based diets. The CP of grass silage, barley, and oats contains considerably less RUP than corn silage and ground corn (NRC, 2001). A smaller contribution of RUP to MP means that the AA composition of RUP has less of an effect on the AA composition of total MP than feeds that are less degradable and have a higher content of RUP in CP. And second, the His content of all of these feeds is low. The His content of barley and oats is lower than the His content of corn (2.3-2.4 vs. 3.1% of CP). The His content of feather meal is considerably lower than the His content of other protein supplements (1.2 vs. 2.0-2.8% of CP). The high content of His in blood (6.4% of CP) is noteworthy and may give blood meal an additional advantage over other protein supplements, particularly when higher forage, lower corn diets are fed.

Ideal Profile of EAA in MP

Based on the above discussion, it seems important that the ideal concentrations of Lys, Met, and His in MP be determined for lactating cows. If these concentrations were known, and diets could be formulated to achieve these ideal concentrations, then the efficiency of use of MP for milk protein production should be maximized. In other words, more milk protein could be produced with the same supply of MP, or the same amount of milk protein could be obtained with less MP.

Both results have been obtained. As reviewed in NRC (2001), and has now been observed on many farms, increasing intestinal supplies of Lys and Met, when they are limiting, increases yields of milk protein. The result is an increased conversion of feed N to milk N. In other cases, it has been shown that milk protein yields can be maintained by feeding less RUP if concentrations of Lys and Met in MP are increased (Noftsker and St-Pierre, 2003; Socha et al., in preparation; farm observations). Again, the result is an increased conversion of feed N to milk N.

Progress has been made in determining the ideal concentrations of Lys and Met in MP. As reviewed at the Four-State Applied Nutrition and Management Conference in 2001 (Schwab, 2001), the NRC (2001) publication contains dose-response plots that relate measured milk protein content and yield responses to changes in predicted percentages of Lys and Met in MP. The plots for milk protein content are shown in Figure 1. The reader is referred to either NRC (2001; p. 81-85) or Schwab (2001) for an explanation of how the plots were generated. It should be noted that the Lys plot was generated from only that portion of the Lys data set where predicted Met was 1.95% or more of MP (solid circles) and that the Met plot was generated from only that portion of the Met data set where predicted Lys was 6.50% or more of MP (solid circles). This was done to help ensure that milk protein responses to supplemental amounts of either AA were not prevented by a deficiency of the other.

The breakpoint estimates for the required concentrations of Lys and Met in MP for maximal content of milk protein were 7.2% and 2.4%, respectively (3.0:1.0 ratio). The breakpoint estimates for the required concentrations of Lys and Met in MP for maximal yield of milk protein were 7.1% and 2.4% (plots not shown). Examination of the dose-response plots indicates little or no expected loss in content or yield of milk protein when Lys and Met in MP are 6.9% and 2.3%, respectively. In fact, our research and field experience indicates no advantage of exceeding these concentrations. Therefore, we consider the latter values (6.9 and 2.3% of MP) to be the absolute highest targets to aim for in diet formulation when using NRC (2001) as the diet evaluation model. However, even these concentrations are difficult to achieve, particularly in high producing cows fed corn based diets. As a result, our “practical recommendations” for percentages of Lys and Met in MP in this case are 6.6 and 2.2, respectively (Figure 2).

For those that use CPM-Dairy or CNCPS as a diet evaluation model, we suggest practical targets of 6.7-6.8% for Lys in MP and 2.2-2.3% for Met.

Balancing for AA in the context of MP and RUP

Based on the above discussion and the senior author’s field experience over the past 2 years, an effective method to balance for Lys and Met involves a 2-step approach.

The first step is to decide what your target values will be for Lys and Met in MP. Will they be 6.6 and 2.2%, or something lower? Whatever you decide, remember two things. First, maintain a Lys:Met ratio in MP of 3.0:1.0 if using NRC (2001) or 3.1:1.0 if using CNCPS or CPM-Dairy. Having a Lys:Met ratio in MP that is either higher (usually the case) or lower (seldom occurs) than 3:1 means that one of the AA is being supplied in excess of need (see Table 2). This has no benefit to the cow. And second, try to get Lys and Met in MP as close to 6.6 and 2.2% as possible. Doing so has the net effect of increasing passage of the

first two limiting AA without the need to increase passage of MP to the small intestine (Table 2). We believe that it is cost effective, even with the current low milk prices, to use the high-Lys protein supplements (soy products and fish and blood meals) in combination with a rumen-protected Met product (RPMet) to get levels of Lys and Met in MP as close to 6.6 and 2.2% as possible (Schwab et al., 2003).

A question that is frequently asked is “what is more important, AA ratios or AA concentrations”? The reason for this question appears to be the result of the frequent emphasis on the 3:1 Lys/Met ratio. In one of the dairy publications about 3 months ago, the following statement was made – “A by-product such as corn gluten meal has a high level of methionine that can be used in combination with blood meal and fish to achieve the desired ratio of 3:1 lysine-methionine”. The author was correct in what he said. However, the emphasis was on AA ratios, with no consideration given to how that strategy affects concentrations of Lys and Met in MP. Purchasing the usually more expensive fish and blood meals only to dilute out their high content of Lys (7.7-9.0% of CP, Table 1) with a low-Lys protein supplement such as corn gluten meal (1.7% of CP) is counter-productive and does not achieve the desired goal of high Lys and Met concentrations in MP. If the decision is made to feed the higher RUP, higher Lys products such as soybean, fish, and blood meals, then the most effective Met supplement is RPMet. Use of a RPMet supplement allows Met in MP to be increased without decreasing Lys.

In a recent paper (Schwab et al., 2003), we made the comparison of achieving a 3:1 Lys-Met ratio in MP using soybean meal, blood meal, and a RPMet product as the primary sources of supplemental AA in a corn based ration vs. using soybean meal, corn distillers grains and corn gluten meal. Diet RDP and RUP concentrations were kept constant between the two diets. In the first case, the NRC (2001) predicted concentrations of Lys and Met in MP were 6.6 and 2.1% (3:1/1.0 ratio) and MP-Lys and Met flows were 191 and 61 g/d. (Note: a 3:1 ratio could have been obtained by including more RPMet in the diet). In the second case, the predicted concentrations of Lys and Met in MP were 5.8 and 1.9% (3.0:1.0 ratio); MP-Lys and Met flows were 170 and 56 g/d.

Field experiences

Presented in Table 3 are the NRC (2001) evaluations of the “before” and “after” diets for six dairy farms in which diet changes were made to increase flows of MP-Lys and Met without increasing flows of total MP. In all cases, Lys concentrations in MP were increased by introducing blood meal and reducing or eliminating distillers grains or a protected soy product. Methionine concentrations in MP were increased to obtain the desired 3:1 Lys-Met ratio in MP by adding a RPMet product (Smartamine M) to the diets. Also presented in Table 3 are the “before” and “after” milk protein and fat percentages. There was no attempt to measure changes in milk yield for most of the herds.

There are three noteworthy observations. First, as evidenced by the high Lys/Met ratios, Met was more limiting than Lys in all of the “before” diets. Therefore, we would conclude that all of the herds would have benefited with higher milk protein concentrations by adding RPMet to the diets. Second, as a result of increasing Lys in MP as well as adding enough RPMet to the diets to achieve a Lys/Met ratio in MP equaled or approximated the desired 3.0/1 ratio, flows of MP-Met were increased 15 to 30%. These are significant increases in the availability of an apparent limiting nutrient and explains why the producers not only observed some rather significant increases in content of milk protein but why in most cases they also thought they observed higher milk yields. And third, in all cases, the diet changes resulted in variable increases in milk fat percentages. This has been an often observed result when Met concentrations in MP are increased (NRC, 2001).

University of New Hampshire Experience

The following discussion describes the “before” and “after” diets for herd 6 and the rationale for the changes that were made. Herd 6 is the University of New Hampshire Research Herd.

For about an 11-month period ending 2 years ago, 75 of our dairy cows were fed a diet containing (DM basis): 29.8% corn silage, 9.6% alfalfa hay, 9.6% grass silage, 15.4% ground corn, 7.4% barley, 4.8% soy hulls, 11.6% soybean meal, 6.4% expeller soybean meal, 1.9% fat, and 3.5% minerals and vitamins. During that period, milk true protein varied between 2.70 and 2.83% and milk fat between 3.4 and 3.7%.

At the end of the 11-month period, the diet was changed. The new diet contained (DM basis): 30.9% corn silage, 12.3% grass silage, 6.0% alfalfa hay, 19.1% corn, 9.4% barley, 3.7% soy hulls, 7.4% soybean meal, 3.7% canola meal, 0.14% urea, 2.2% of a highly digestible animal protein blend (Venture Milling) that contained 0.075% Smartamine M and 0.10% Rhodimet AT-88, 1.9% fat, and 3.1% minerals and vitamins. The animal protein blend replaced the expeller soybean meal to increase Lys in MP. Smartamine M was added to achieve the desired 3.0/1.0 Lys to Met ratio in MP. The canola meal and urea replaced some of the soybean meal to provide a more diverse mix of RDP and to lower the cost of RDP. And finally, RDP and RUP were decreased to eliminate some of what NRC (2001) indicated to be a surplus and to offset the higher cost of the animal protein blend. The new diet contained 17.2% CP as compared to 18.1% for the old diet. According to NRC (2001), the new diet contained 10.6% RDP (instead of 10.8%) and 6.6% RUP (instead of 7.3%). Because of the decrease in RUP, predicted MP flows to the small intestine were decreased from 3071 to 2809 g/d (8% reduction). RDP was lowered by 2%. However, predicted concentrations of Lys and Met in MP increased from 6.34% and 1.73% to 6.55% and 2.20%, respectively. Therefore, even though predicted passage of MP was 8% less, the

predicted flow of MP-Met (previously the “weakest link”) was increased from 53 to 61 g/d, a 16% increase. Predicted MP-Lys flows decreased from 195 to 184 g/d. This was not considered to be a problem because amounts greater than what would be needed to achieve a Lys/Met ratio of 3.0/1.0 using NRC (2001) would be considered to be a surplus.

The cows were switched gradually over a 10-d period to the new diet. For the 2-wk period preceding the transition to the new diet, milk protein concentrations averaged 2.82%. Although considered to be low, this level of milk protein was at the high end of the range (2.70% to 2.83%) for the preceding 11-month period. One week after the change, milk protein concentrations had increased to 3.01%. At the end of wk 2, protein increased to 3.06% and by wk 4 it had increased to 3.13%. Thereafter, and for the next couple of months while it was being monitored, milk protein stabilized between 3.12 and 3.16%. As expected because of the decrease in ration CP, milk urea N decreased from an average of 14.5 to an average of 12.4 mg/dL. Milk fat concentrations also increased. It appeared that milk yields increased but it was difficult to determine that as the cows (on average) were advancing in days-in-milk.

The economics of the diet changes were most favorable. Because we assumed that milk yield was not increased, we also assumed that DM intake was not affected. A cost analysis of the diet indicated an approximate 5 cents per cow per day increase in feed costs. However, because of the increases in milk protein and milk fat concentrations, milk income was increased by \$0.70 per cow per day. The increased income-over-feed-costs (IOFC) was \$0.65 (assuming no increase in milk yield or feed intake).

Predicting Production from MP-Lys and MP-Met

Is balancing for Lys and Met in MP always profitable? The use of high quality protein and RPMet supplements to increase intestinal flows of MP-Lys and Met continue to be questioned as to their economic value. Do we keep the high-Lys, high-digestible blood, fish and soybean meals and the protected Met supplements in the diets at our normal inclusion levels, or just feed less? Or, do we let cost per ton of feed take center stage and scrap these feeds? Unfortunately, the current diet evaluation models that predict passage of MP-AA to the small intestine (e.g., NRC, 2001; CNCPS, and CPM-Dairy) in their present form are not useful in predicting the effect that changes in supplies of MP-Lys and MP-Met have on milk and milk component production. Until such systems are in place, it will remain difficult to predict the effect that changes in protein supplementation strategies have on milk and milk protein yields and the resulting effects on IOFC.

Schwab et al. (2003) used the NRC (2001) model in conjunction with published experiments to examine the relationships between predicted supplies of MP, MP-Met, and MP-Lys and yields of milk and milk protein. That effort has been extended for this paper.

Over 300 diets from experiments published in the Journal of Dairy Science were entered into the NRC (2001) model. In most of these experiments the objective was to compare the effects of feeding different protein supplements on milk production and milk composition, and in some cases, passage of N fractions to the small intestine. Relevant data from the Summary and Duodenal Amino Acid Supply Reports were recorded.

To generate plots of measured yields of milk and milk protein vs. predicted supplies of MP, data were restricted to diets in which NE-allowable milk was higher than MP-allowable milk, and actual milk yield was between minus 6 kg and plus 6 kg of MP-allowable milk. The former restriction was imposed to help ensure that MP was more limiting than NE. The latter restriction was imposed to avoid the use of experiments in which factors other than MP or NE limited lactation performance or situations where excessive protein mobilization may have been occurring.

To generate plots of measured yields of milk and milk protein vs. predicted supplies of MP-Lys and MP-Met, data were restricted to diets in which MP balance was within -250 and +100 g/d of zero balance. This was done with the hope of further ensuring that Lys and Met were limiting. For the Met plots, we imposed the restriction that the Lys/Met ratio in MP had to be greater than 3.0/1.0 to make more certain that Met was more limiting than Lys. For the Lys plots, we wanted to add the restriction that the ratio of Lys to Met in MP had to be less than 3.0/1.0 to ensure that Lys was more limiting in MP than Met. However, only in a few cases was the ratio of Lys to Met in MP less than 3.0/1.0. Therefore, to give ourselves an adequate number of data points from which to get some idea of the relationship between yields of milk and milk protein vs. predicted supplies of Lys, diets yielding predicted Lys/Met ratios up to 3.25/1.0 were used.

The resulting plots are presented in Figure 3. There are at least three observations that are worthy of mention. First, in all cases (for MP, MP-Met, and MP-Lys), it appears that protein yields can be predicted more accurately than milk yields. This would be expected because of the changes in milk protein percentages that often occur with changes in protein nutrition. Second, as expected, predicting yields of milk and milk protein from intestinal supplies of the most limiting AA is more precise than predicting yields from MP supply. Third, and while the current data is too limited and not adequate for this exercise, it appears that a very strong relationship exists between milk and milk protein yields and predicted MP-Lys supplies. This should probably be expected given the fact that Lys, unlike Met, has only one function in the body, i.e., protein synthesis.

Conclusions

Balancing diets to optimize Lys and Met nutrition is important to maximizing milk and milk protein yields. As expected, it appears that establishing relationships between predicted supplies of the most limiting AA in the diet and milk and milk protein yields will allow for more accurate prediction of changes in milk protein production when changes in protein nutrition are made. Of concern is the lack of a rumen-protected Lys product and therefore, the inability to achieve desired concentrations of Lys in MP when high corn diets are fed.

References

- Huhtanen, P., V. Vanhatalo, and T. Varvikko. 2002. Effects of abomasal infusions of histidine, glucose, and leucine on milk production and plasma metabolites of dairy cows fed grass silage diets. *J. Dairy Sci.* 85:204-216.
- Kim, C. H., T. G. Kim, J. J. Choung, and D. G. Chamberlain. 1999. Determination of the first limiting amino acid for milk production in dairy cows consuming a diet of grass silage and a cereal-based supplement containing feather meal. *J. Sci. Food Agric.* 79:1703-1708.
- Kim, C. H., T. G. Kim, J. J. Choung, and D. G. Chamberlain. 2000. Variability in the ranking of the three most-limiting amino acids for milk protein production in dairy cows consuming grass silage and a cereal-based supplement containing feather meal. *J. Sci. Food Agric.* 80:1386-1392.
- Kim, C. H., T. G. Kim, J. J. Choung, and D. G. Chamberlain. 2001a. Effects of intravenous infusion of amino acids and glucose on the yield and concentration of milk protein in dairy cows. *J. Dairy Res.* 68:27-34.
- Kim, C. H., T. G. Kim, J. J. Choung, and D. G. Chamberlain. 2001b. Estimates of the efficiency of transfer of L-histidine from blood to milk when it is the first-limiting amino acid for secretion of milk protein in the dairy cow. *J. Sci. Food Agric.* 81:1150-1155.
- Korhonen, M., A. Vanhatalo, T. Varvikko, and P. Huhtanen. 2000. Responses to graded doses of histidine in dairy cows fed grass silage diets. *J. Dairy Sci.* 83:2596-2608.
- National Research Council. 2001. *Nutrient Requirements of Dairy Cattle*. 7th rev. ed. Natl. Acad. Sci., Washington, DC.
- National Research Council. 1998. *Nutrient Requirements of Swine*. 10th rev. ed. Natl. Acad. Sci., Washington, DC.
- National Research Council. 1994. *Nutrient Requirements of Poultry*. 9th rev. ed. Natl. Acad. Sci., Washington, DC.

- National Research Council. 1989. Nutrient Requirements of Dairy Cattle. 7th rev. ed. Natl. Acad. Sci., Washington, DC.
- National Research Council. 1978. Nutrient Requirements of Dairy Cattle. 5th rev. ed. Natl. Acad. Sci., Washington, DC.
- National Research Council. 1971. Nutrient Requirements of Dairy Cattle. 4th rev. ed. Natl. Acad. Sci., Washington, DC.
- Noftsgger, S. and N. R. St-Pierre. 2003. Supplementation of Met and selection of highly digestible rumen undegradable protein to improve nitrogen efficiency for milk production. *J. Dairy Sci.* 86: 958-969.
- O'Connor, J. D., C. J. Sniffen, D. G. Fox, and W. Chalupa. 1993. A net carbohydrate and protein system for evaluating cattle diets: IV. Predicting amino acid adequacy. *J. Anim. Sci.* 71:1298-1311.
- Schwab, C. G. 2001. Dairy NRC 2001: Protein System. In: Proc. Four-State Applied Nutrition and Management Conference, La Crosse, WI, p. 25-34.
- Schwab, C. G., R. S. Ordway, and N. L. Whitehouse. 2003. The latest on amino acid feeding. In: Proc. Southwest Nutrition and Management Conference, Phoenix, AZ, p. 27-41.
- Socha, M. T., D. E. Putnam, B. D. Garthwaite, N. L. Whitehouse, N. A. Kierstead, C. G. Schwab, G. A. Ducharme, and J. C. Robert. 2003. Supplementing diets of prepartum and early lactation cows with rumen-stable amino acids. *J. Dairy Sci.* (In preparation).
- Vanhatalo, A., P. Huhtanen, V. Toivonen, and T. Varvikko. 1999. Response of dairy cows fed grass silage diets to abomasal infusions of histidine alone or in combinations with methionine and lysine. *J. Dairy Sci.* 82:2674-2685.

Table 1. A comparison of the essential amino acid composition of body lean tissue, milk, and ruminal bacteria with that of some common feeds¹.

Item	Arg	His	Ile	Leu	Lys	Met	Phe	Thr	Trp	Val
	----- (% of CP) -----									
Lean tissue	6.6	2.5	2.8	6.7	6.4	2.0	3.5	3.9	0.6	4.0
Milk	3.4	2.7	5.8	9.2	7.6	2.7	4.8	3.7	1.5	5.9
Bacteria	5.1	2.0	5.7	8.1	7.9	2.6	5.1	5.8	-	6.2
Alfalfa silage	3.9	1.7	3.9	6.4	4.4	1.4	4.2	3.8	0.9	5.0
Corn silage	2.0	1.8	3.3	8.6	2.5	1.5	3.8	3.2	0.4	4.5
Grass silage	3.1	1.7	3.6	6.1	3.3	1.2	4.4	3.3	1.1	4.9
Barley	5.1	2.3	3.5	7.0	3.6	1.7	5.1	3.4	1.2	4.9
Corn	4.6	3.1	3.3	11.2	2.8	2.1	4.6	3.6	0.7	4.0
Oats	6.8	2.4	3.8	7.3	4.2	2.9	5.2	3.5	1.2	5.2
Wheat	4.7	2.4	3.3	6.6	2.8	1.6	4.6	2.9	1.2	4.2
Brewers grains	5.8	2.0	3.9	7.9	4.1	1.7	4.6	3.6	1.0	4.8
Canola meal	7.0	2.8	3.8	6.8	5.6	1.9	4.1	4.4	1.5	4.7
Corn DDG w/sol	4.1	2.5	3.7	9.6	2.2	1.8	4.9	3.4	0.9	4.7
Corn gluten meal	3.2	2.1	4.1	16.8	1.7	2.4	6.4	3.4	0.5	4.6
Cottonseed meal	11.1	2.8	3.1	5.9	4.1	1.6	5.3	3.2	1.2	4.2
Soybean meal	7.3	2.8	4.6	7.8	6.3	1.4	5.3	4.0	1.3	4.6
Sunflower meal	8.2	2.6	4.1	6.4	3.6	2.3	4.6	3.7	1.2	5.0
Blood meal	4.4	6.4	1.3	12.8	9.0	1.2	6.9	4.3	1.6	8.7
Feather meal	6.9	1.2	4.9	8.5	2.6	0.8	4.9	4.7	0.7	7.5
Fish meal	5.8	2.8	4.1	7.2	7.7	2.8	4.0	4.2	1.1	4.8
Meat meal	7.1	2.1	3.0	6.3	5.4	1.4	3.6	3.4	0.7	4.4

¹ Amino acid values for lean tissue, milk, and ruminal bacteria are from O'Connor et al. (1993) and amino acid values for feeds are from NRC (2001).

Table 2. Effect of NRC (2001) predicted percentages of Lys and Met in MP on calculated flows of MP-Lys and MP-Met, and the amounts of MP-Lys and Met that can be used for protein synthesis.

Predicted Lys/Met in MP (%)	Lys/Met ratio	Flows ¹		Used for protein synthesis ²	
		MP-Lys	MP-Met	MP-Lys	MP-Met
		----- (g/d) -----			
6.4/1.7	3.8/1	179	48	144	48
5.8/1.7	3.4/1	162	48	144	48
5.7/1.9	3.0/1	160	53	159	53
5.8/2.1	2.8/1	162	59	162	54
6.3/2.0	3.2/1	176	56	168	56
6.6/2.2	3.0/1	185	62	186	62

¹ Calculations are based on a predicted MP supply of 2,800 g/d.

² Based on the assumption that the optimum Lys/Met ratio in MP is 3:1 and the understanding that any AA supplied in excess of need for protein synthesis is not used for protein synthesis and therefore, is catabolized and used for energy.

Table 3. NRC (2001) evaluations of “Before” and “After” diets for six commercial farms in which concentrations of Met and Lys in MP were increased.

Item	Herd 1		Herd 2		Herd 3	
	Before	After	Before	After	Before	After
CP, %	17.7	17.8	18.3	18.6	18.3	18.1
RDP, %	10.9	11.2	11.0	11.2	11.4	11.3
RUP, %	6.8	6.6	7.3	7.4	6.9	6.8
MP, g/d	3054	2984	3159	3131	3062	3040
Lys, %MP	5.78	6.53	5.74	6.20	5.84	6.18
Met, %MP	1.65	2.17	1.68	2.08	1.68	2.01
MP-Lys, g/d	177	195	181	194	179	188
MP-Met, g/d	50	65	53	65	51	61
Lys/Met	3.5/1	3.0/1	3.4/1	3.0/1	3.5/1	3.1/1
Milk protein, %	3.06	3.33	2.99	3.12	3.02	3.22
Milk fat, %	3.81	3.92	3.56	3.66	3.61	3.72

Item	Herd 4		Herd 5		Herd 6	
	Before	After	Before	After	Before	After
CP, %	19.1	18.2	17.6	17.0	18.1	17.2
RDP, %	12.0	11.2	10.4	10.3	10.8	10.6
RUP, %	7.1	6.9	7.2	6.7	7.3	6.6
MP, g/d	3107	3030	3073	3035	3071	2809
Lys, %MP	5.84	6.25	6.34	6.76	6.37	6.55
Met, %MP	1.67	2.04	1.73	2.35	1.73	2.20
MP-Lys, g/d	182	189	195	193	184	174
MP-Met, g/d	52	62	53	61	50	58
Lys/Met	3.5/1	3.0/1	3.7/1	3.2/1	3.7/1	3.0/1
Milk protein, %	3.00	3.20	3.20	3.50	2.82	3.16
Milk fat, %	3.49	3.64	3.90	4.30	3.32	3.78





Figure 1. Milk protein content responses as a function of percent Lys and Met in MP. For the Lys plot, the regression analysis was limited to data where Met was predicted to be 1.95% or more of MP (solid circles). For the Met plot, the regression analysis was limited to data where Met was predicted to be 6.50% or more of MP (solid circles) (NRC, 2001).

Figure 2. Optimum versus practical levels of Lys and Met in metabolizable protein.

